

Product Datasheet **GenieFluor Red 780 Anti-Human HLA-DR Antibody [L243]** Catalogue Code: AGEL2980

Antibody Data

Product SKU:	AGEL2980	Clone:	L243
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	HLA class II histocompatibility antigen DR alpha/ DRB1-15 beta chain;MHC class II antigen DRA;HLA-DRA1/DRB1;DRA/DRB1; P01903 P01911 HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells.		
Form:	Liquid	780 Excitation and Emission Spectra	
Conjugation:	Genie FluorRed 780	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 -	
Host Species:	Mouse	(%) 60	
Isotype:	Mouse IgG2a, к	$\mathbf{E}_{20}^{0} \underbrace{\begin{array}{c} 40 \\ 0 \\ 350 \end{array}}_{350 \hspace{0.5cm} 400 \hspace{0.5cm} 450 \hspace{0.5cm} 500 \hspace{0.5cm} 550 \hspace{0.5cm} 600 \hspace{0.5cm} 650 \hspace{0.5cm} 700 \hspace{0.5cm} 750 \hspace{0.5cm} 800 \hspace{0.5cm} 850 \hspace{0.5cm} 850 \hspace{0.5cm} Wavelength (nm) \end{array}}$	

Isotype Control: Genie Fluor Red 780 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL2980]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.