

#### **Product Datasheet**

# GenieFluor Red 780 Anti-Human CD14 Antibody [M5E2]

Catalogue Code: AGEL2998

### Antibody Data

Product SKU: AGEL2998 Clone: M5E2

Applications: FCM

Reactivity: Human

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information:**

Alternate Names: Monocyte differentiation antigen CD14;CD14;Myeloid cell-specific leucine-rich

glycoprotein;

Uniprot ID: P08571

**Background**: CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein also

known as LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens, and in the upregulation of adhesion molecules

and expression of cytokines in monocytes and neutrophils.


Form: Liquid

Conjugation: Genie FluorRed 780

Size: 20 Tests, 100 Tests, 200 Tests

Host Species: Mouse

**Isotype:** Mouse IgG2a, κ



Isotype Control: Genie Fluor Red 780 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL2998]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.