

Product Datasheet

GenieFluor Violet 450 Anti-Human CD45RA Antibody [HI100] Catalogue Code: AGEL3003

Antibody Data

Product SKU:	AGEL3003	Clone:	HI100	
Applications:	FCM			
Reactivity:	Human			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

	Descrites time transis a mattein absorb store O.D.target, OA.L., 5.T000.OD.45.		
Alternate Names:	Receptor-type tyrosine-protein phosphatase C;Ptprc;L-CA;Ly-5;T200;CD45;		
Uniprot ID:	P08575		
Background:	CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines.		
Form:	Liquid	450 Excitation and Emission Spectra	
Conjugation:	Genie FluorViolet 450	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse		
Isotype:	Mouse IgG2b, κ	20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	
Isotype Control:	Genie Fluor Violet 450 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL3003]		

Isotype Control: Genie Fluor Violet 450 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL3003]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.