

Product Datasheet GenieFluor Red 780 Anti-Mouse CD25 Antibody [PC-61.5.3]

Catalogue Code: AGEL3012

## Antibody Data

Product SKU:	AGEL3012	Clone:	PC-61.5.3	
Applications:	FCM			
Reactivity:	Mouse			

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID: Background:	Interleukin-2 receptor subunit alpha;IL2RA;IL-2 receptor subunit alpha;IL-2-RA;IL-2R subunit alpha;IL2-RA;TAC antigen;p55;p55; P01590 CD25 is a 55 kD glycoprotein, also known as the low affinity IL-2Rα, Ly-43, p55, or Tac. It is expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2Rβ) and CD132(common γ chain), CD25 forms the high affinity signaling IL-2 receptor.		
Form:	Liquid	'80 Excitation and Emission Spectra	
Conjugation:	Genie FluorRed 780	100 -	
Size:	50 Tests, 100 Tests, 200 Tests	80 -	
Host Species:	Rat	(%) po 00	
Isotype:	Rat IgG1, κ	$\frac{1}{9} \begin{bmatrix} 40 \\ 20 \\ 0 \\ 350 \end{bmatrix} = \frac{1}{400} \begin{bmatrix} 20 \\ 450 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} 20 \\ 500 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} 20 \\ 500 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} 20 \\ 500 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} 20 \\ 700 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} 20 \\ 800 \end{bmatrix} = \frac{1}{500} \begin{bmatrix} $	
Isotype Control:	Genie Fluor Red 780 Rat IgG1, κ Isotype Control[HRPN] [Product AGEL3012]		

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.