

Product Datasheet

GenieFluor Red 780 Anti-Human/Mouse CD44 Antibody [IM7]

Catalogue Code: AGEL3069

Antibody Data

Product SKU: AGEL3069 Clone: IM7

Applications: FCM

Reactivity: Human; Mouse

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: CD44 antigen; CD44; CDw44; Epican; Phagocytic glycoprotein 1; PGP-1; Phagocytic

glycoprotein I;PGP-I;CD44;LHR; MDU2; MDU3; MIC4;

Uniprot ID: P15379 P16070

Background: CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is

expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte

aggregation.


Form: Liquid

Conjugation: Genie FluorRed 780

Size: 25µg, 100µg

Host Species: Rat

Isotype: Rat IgG2b, κ



Isotype Control: Genie Fluor Red 780 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL3069]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/106 cells in 100 μ L volume].