

**GenieFluor Red 780 Anti-Mouse CD80 Antibody [16-10A1]**  
 Catalogue Code: AGEL3114

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL3114</b>	<b>Clone:</b>	<b>16-10A1</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** T-lymphocyte activation antigen CD80;Cd80;Activation B7-1 antigen;B7;

**Uniprot ID:** Q00609

**Background:** CD80 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD80 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD80 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD80 appears to be expressed later in the immune response than CD86. CD80 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.


**Form:** Liquid

**Conjugation:** Genie FluorRed 780

**Size:** 25µg, 100µg

**Host Species:** Armenian Hamster

**Isotype:** Armenian Hamster IgG



**Isotype Control:** Genie Fluor Red 780 Armenian Hamster IgG Isotype Control[PIP] [Product AGEL3114]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10<sup>6</sup> cells in 100 µL volume].

---