

Product Datasheet **PE/GenieFluor 594 Anti-Human CD62L Antibody [DREG56]** Catalogue Code: AGEL3145

Antibody Data

Product SKU:	AGEL3145	Clone:	DREG56
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	 L-selectin;Sell;CD62 antigen-like family member L;LAM-1;LECAM1;Lymph node homing receptor;Ly-22; CD62L;Lnhr;Ly22; P14151 CD62L is a 74-95 kD single chain type I glycoprotein referred to as L-selectin or LECAM-1. It is expressed on most peripheral blood B cells, subsets of T and NK cells, monocytes, granulocytes, and certain hematopoietic malignant cells. CD62L binds to carbohydrates present on certain glycoforms of CD34, glycam-1, and MAdCAM-1 and with a low affinity to anionic oligosaccharide sequences related to sialylated Lewis X (sLex, CD15s) through its C-type lectin domain. CD62L is important for the homing of naïve lymphocytes to high endothelial venules in peripheral lymph nodes and Peyer's patches. It also plays a role in leukocyte rolling on activated endothelial cells. 		
Form:	Liquid	594 Excitation and Emission Spectra	
Conjugation:	PE/Genie Fluor594	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 -	
Host Species:	Mouse	(%) 00 - 00 - 00 - 00 - 00 - 00 - 00 - 00	
Isotype:	Mouse IgG1, κ	$\frac{1}{20} = \frac{1}{350} = \frac{1}{400} = \frac{1}{450} = \frac{1}{500} = \frac{1}{550} = \frac{1}{600} = \frac{1}{650} = \frac{1}{700} = \frac{1}{750} = \frac{1}{800} = \frac{1}{850}$ Wavelength (nm)	
Isotype Control:	PE/Genie Fluor 594 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3145]		

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.