

#### **Product Datasheet**

# PerCP/Cyanine5.5 Anti-Human CD34 Antibody [581]

Catalogue Code: AGEL3165

## Antibody Data

Product SKU: AGEL3165 Clone: 581

Applications: FCM

Reactivity: Human

### **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Gp105-120; My10;

Uniprot ID: P28906

Background: CD34, also known as gp105-120, is a type I monomeric sialomucin-like

glycophosphoprotein with an approximate molecular weight of 105-120 kD. Selectively expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nervous tissue, CD34 is a commonly used marker to identify human hematopoietic stem/progenitor cells. According to the differential sensitivity to enzymatic cleavage, four groups of epitopes of CD34 have been described. CD34 mediates cell adhesion and lymphocytes homing

through binding to L-selectin and E-selectin ligands.


Form: Liquid

**Conjugation:** PerCP/Cyanine 5.5

Size: 20 Tests, 100 Tests, 200 Tests

Host Species: Mouse

**Isotype:** Mouse IgG1, κ



Ex:440;480;675 nm; Em:675 nm

**Isotype Control:** PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3165]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



Stability & Storage:

Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.