

Product Datasheet **PE/GenieFluor 594 Anti-Human CD45RA Antibody [HI100]** Catalogue Code: AGEL3211

Antibody Data

| Product SKU: | AGEL3211 | Clone: | HI100 |
|---------------|----------|--------|-------|
| Applications: | FCM | | |
| Reactivity: | Human | | |

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

| Alternate Names: | Receptor-type tyrosine-protein phosphatase C;Ptprc;L-CA;Ly-5;T200;CD45; | | |
|------------------|--|--|--|
| Uniprot ID: | P08575 | | |
| Background: | CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines. | | |
| Form: | Liquid | 94 Excitation and Emission Spectra | |
| Conjugation: | PE/Genie Fluor594 | 100 - | |
| Size: | 20 Tests, 100 Tests, 200 Tests | 80 - | |
| Host Species: | Mouse | | |
| Isotype: | Mouse IgG2b, κ | $\mathbf{F}_{2}^{4_{0}} = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0$ | |
| Isotype Control: | PE/Genie Fluor 594 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL3211] | | |

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.