

#### **Product Datasheet**

# PE Anti-Human/Mouse IL-5 Antibody [TRFK5]

Catalogue Code: AGEL3280

# Antibody Data

Product SKU: AGEL3280 Clone: TRFK5

Applications: ICFCM

Reactivity: Human; Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information:**

**Alternate Names:** Interleukin-5;IL-5;B-cell differentiation factor I;T-cell replacing factor;TRF;

**Uniprot ID:** P05113 P04401

**Background**: IL-5 is a homodimeric, disulphide-linked protein produced by T-cells. Monomeric human

IL-5 is a 126 amino acid protein with a reported molecular weight of 26 kD for the homodimeric protein. Mouse and human IL-5 are approximately 70% identical. IL-5 has been shown to promote the growth of immature hematopoietic BFU-E progenitors, stimulate the activation and differentiation of eosinophils, and promote the generation of cytotoxic lymphocytes. Mouse IL-5 induces the differentiation and proliferation of preactivated B-cells and stimulates the production and secretion of IgM and IgA by B-cells stimulated with bacterial endotoxin. The TRFK5 antibody can neutralize the bioactivity of

natural or recombinant IL-5.


Form: Liquid

Conjugation: PE

Size: 20 Tests, 100 Tests, 200 Tests

Host Species: Rat

**Isotype:** Rat IgG1, κ



Ex:495;565 nm; Em:575 nm

**Isotype Control**: PE Rat IgG1, κ Isotype Control[HRPN] [Product AGEL3280]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



Stability & Storage:

Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.