

Product Datasheet **GenieFluor Red 780 Anti-Human CD56 Antibody [5.1H11]** Catalogue Code: AGEL3285

Antibody Data

Product SKU:	AGEL3285	Clone:	5.1H11
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Neural cell adhesion molecule 1;NCAM1;N-CAM-1;NCAM-1;NCAM;		
Uniprot ID: Background:	P13591 CD56 is a single transmembrane glycoprotein also known as NCAM (Neural Cell Adhesion		
	Molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is expressed on NK cells and NK-T cells. CD56 is also expressed in the brain (cerebellum and cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparin sulfate.		
Form:	Liquid	780 Excitation and Emission Spectra	
Conjugation:	Genie FluorRed 780	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 2 60	
Host Species:	Mouse	(%) 60	
Isotype:	Mouse IgG1, κ	$\frac{5}{20} \frac{40}{0} \frac{1}{350} \frac{1}{400} \frac{1}{450} \frac{1}{500} \frac{1}{550} \frac{1}{600} \frac{1}{650} \frac{1}{700} \frac{1}{750} \frac{1}{800} \frac{1}{850} \frac{1}{850} \frac{1}{100} \frac{1}{1$	

Isotype Control: Genie Fluor Red 780 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3285]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.