

Product Datasheet

GenieFluor 488 Anti-Human CD68 Antibody [Y1/82A] Catalogue Code: AGEL3313

Antibody Data

Product SKU:	AGEL3313	Clone:	Y1/82A
Applications:	ICFCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

D24040		
P34810		
CD68 is a 110 kD glycoprotein, also known as macrosialin, belonging to the sialomucin family. It is closely related to the family of acidic, highly glycosylated lysosomal-associated membrane proteins (LAMPs). CD68 is predominately expressed in cytoplasmic granules of monocytes/macrophages, dendritic cells, and granulocytes. It is one of the useful myeloid cell markers. Further studies have shown that CD68 is also expressed by a subset of hematopoietic progenitors, γ/δ T cells, NK cells, LAK cells, subset of B cells, fibroblasts, and endothelial cells. The biological function of CD68 is still unknown.		
Liquid	488 Excitation and Emission Spectra	
Genie Fluor488	100	
20 Tests, 100 Tests, 200 Tests	80 - E	
Mouse		
Mouse IgG2b, κ	E 40 20 0 350 400 450 550 600 650 700	
	CD68 is a 110 kD glycoprotein, family. It is closely related to the f membrane proteins (LAMPs). CD of monocytes/macrophages, den myeloid cell markers. Further stud of hematopoietic progenitors, γ/δ and endothelial cells. The biologi Liquid Genie Fluor488 20 Tests, 100 Tests, 200 Tests Mouse Mouse IgG2b, κ	

Isotype Control: Genie Fluor 488 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL3313]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.