



Product Datasheet

Purified Anti-Human CD16 Antibody [3G8]

Catalogue Code: AGEL3348

Antibody Data

Product SKU:	AGEL3348	Clone:	3G8
Applications:	FCM;Block		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	FcγRIII;Fc gamma receptor;Fc gamma receptor 3;
Uniprot ID:	P08637 O75015
Background:	CD16 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD16 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD16 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD16 appears to be expressed later in the immune response than CD86. CD16 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.
Form:	Liquid
Conjugation:	Unconjugated
Size:	25µg, 100µg
Host Species:	Mouse
Isotype:	Mouse IgG1, κ
Isotype Control:	Purified Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3348]
Storage Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
Shipping:	Biological ice pack at 4°C

Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5 \mu\text{g}$ per 10^6 cells in $100 \mu\text{L}$ volume or $100 \mu\text{L}$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
