TRAF1 Rabbit Polyclonal Antibody

CAB0150

Size:

Product Information

20uL, 50uL, 100uL, 200uL




The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor

(TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from

various receptors of the TNFR superfamily. This protein and TRAF2 form a heterodimeric complex, which is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. **Observed MW:** The protein complex formed by this protein and TRAF2 also interacts with inhibitor-ofapoptosis proteins (IAPs), and thus mediates the anti-apoptotic signals from TNF receptors. 50kDa The expression of this protein can be induced by Epstein-Barr virus (EBV). EBV infection **Calculated MW:** membrane protein 1 (LMP1) is found to interact with this and other TRAF proteins; this interaction is thought to link LMP1-mediated B lymphocyte transformation to the signal 33kDa/46kDa transduction from TNFR family receptors. Three transcript variants encoding two different isoforms have been found for this gene. **Applications:** Immunogen information WB Gene ID: **Reactivity:** 7185 Human, Mouse, Rat Uniprot Q13077 **Antibody Information** Synonyms: **Recommended dilutions:** TRAF1; EBI6; MGC:10353 WB 1:500 - 1:2000 Source: Rabbit Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 217-416 of human TRAF1 (NP_005649.1). **Isotype:** lgG Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. **Purification:** Affinity purification

Protein Background



Western blot analysis of extracts of various cell lines, using TRAF1 antibody (CAB0150) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.