

BRCA1 Rabbit Polyclonal Antibody



CAB0212

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

Refer to figures

Calculated MW:

7kDa/78-85kDa/202-210kDa

Applications:

WB IHC IF

Reactivity:

Human, Mouse

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50
- 1:200 IF 1:20 - 1:50

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Protein Background

This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast and ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiological function of this gene. Many alternatively spliced transcript variants, some of which are disease-associated mutations, have been described for this gene, but the full-length nature of only some of these variants has been described. A related pseudogene, which is also located on chromosome 17, has been identified.

Immunogen information

Gene ID:

672

Uniprot

P38398

Synonyms:

BRCA1; BRCAI; BRCC1; BROVCA1; FANCS; IRIS; PNCA4; PPP1R53;
PSCP; RNF53


Immunogen:

A synthetic peptide corresponding to a sequence within amino acids 120-220 of human BRCA1 (NP_009225.1).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images



Immunohistochemistry of paraffin-embedded mouse brain using BRCA1 antibody (CAB0212) at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human breast cancer using BRCA1 antibody (CAB0212) at dilution of 1:100 (40x lens).



Immunofluorescence analysis of HeLa cells using BRCA1 Rabbit pAb (CAB0212) at dilution of 1:100. Blue: DAPI for nuclear staining.