

## CAB0217

---

**Product Information**

<b>Product SKU:</b>	CAB0217	<b>Gene ID:</b>	100506742	<b>Size:</b>	20uL, 100uL
<b>Clone No:</b>	-	<b>Host Species:</b>	Rabbit	<b>Reactivity:</b>	Human,Mouse,Rat

---

**Additional Information**

<b>Observed MW:</b>	55kDa	<b>Conjugate:</b>	Unconjugated
<b>Calculated MW:</b>	39kDa	<b>Isotype:</b>	IgG

---

**Immunogen Information**

<b>Background:</b>	Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their substrate molecules. This gene is most highly related to members of the ICE subfamily of caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates apoptosis in response to endoplasmic reticulum stress. However, in humans this gene contains a polymorphism for the presence or absence of a premature stop codon. The majority of human individuals have the premature stop codon and produce a truncated non-functional protein. The read-through codon occurs primarily in individuals of African descent and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe sepsis. Several alternatively spliced transcript variants have been noted for this gene.
<b>Recommended Dilution:</b>	WB,1:500 - 1:1000 IHC-P,1:50 - 1:200
<b>Synonyms:</b>	CASP-12; CASP12P1; Caspase-12
<b>Purification Method:</b>	Affinity purification
<b>Immunogen:</b>	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human Caspase-12 (NP_001177945.2).
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.