## **IRF4 Rabbit Polyclonal Antibody**



## CAB0524

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

52kDa

Calculated MW:

51kDa

**Applications:** 

WB

Reactivity:

Human, Mouse, Rat

**Protein Background** 

The protein encoded by this gene belongs to the IRF (interferon regulatory factor) family of transcription factors, characterized by an unique tryptophan pentad repeat DNA-binding domain. The IRFs are important in the regulation of interferons in response to infection by virus, and in the regulation of interferon-inducible genes. This family member is lymphocyte specific and negatively regulates Toll-like-receptor (TLR) signaling that is central to the activation of innate and adaptive immune systems. A chromosomal translocation involving this gene and the IgH locus, t(6;14)(p25;q32), may be a cause of multiple myeloma. Alternatively spliced transcript variants have been found for this gene.

Immunogen information

Gene ID: 3662

Uniprot Q15306

Synonyms:

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source: Rabbit

Immunogen:

Recombinant fusion protein containing a sequence corresponding

to amino acids 150-350 of human IRF4 (NP\_002451.2).

IRF4; LSIRF; MUM1; NF-EM5; SHEP8; IRF-4

Storage: Isotype:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% IgG

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using IRF4 antibody (CAB0524) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.