## **PRKACA Rabbit Polyclonal Antibody**



## **CAB0798**

**Product Information** 

Size:

100uL, 200uL

**Observed MW:** 

41kDa

**Calculated MW:** 

39kDa/40kDa

**Applications:** 

**WB IHC** 

Reactivity:

Human, Mouse

**Antibody Information** 

**Recommended dilutions:** WB 1:500 - 1:2000 IHC 1:50

- 1:200

Source:

Isotype:

IgG

Rabbit

**Protein Background** 

This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. Tissue-specific isoforms that differ at the N-terminus have been described, and these isoforms may differ in the post-translational modifications that occur at the N-terminus of some isoforms.

Immunogen information

Gene ID:

5566

Uniprot

P17612

Synonyms:

PRKACA; PKACA; PPNAD4

Immunogen:


A synthetic peptide of human PRKACA

Storage:

Purification: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

Affinity purification sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extracts of HeLa cells, using PRKACA antibody (CAB0798). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.