

NRXN1 Rabbit Polyclonal Antibody



CAB10066

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

162kDa

Calculated MW:

15kDa/46kDa/161kDa/164kDa/169kDa

Applications:

WB

Reactivity:

Human, Mouse, Rat

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Protein Background

This gene encodes a single-pass type I membrane protein that belongs to the neurexin family. Neurexins are cell-surface receptors that bind neuroligins to form Ca²⁺-dependent neurexin/neuroligin complexes at synapses in the central nervous system. This complex is required for efficient neurotransmission, and is involved in the formation of synaptic contacts. Three members of this gene family have been studied in detail and are estimated to generate over 3,000 variants through the use of two alternative promoters (alpha and beta) and extensive alternative splicing in each family member. Recently, a third promoter (gamma) was identified for this gene in the 3' region. The RefSeq project has decided to create only a few representative transcript variants of the multitude that are possible. Mutations in this gene are associated with Pitt-Hopkins-like syndrome-2.

Immunogen information

Gene ID:

9378

Uniprot

P58400/Q9ULB1

Synonyms:

NRXN1; Hs.22998; PTHSL2; SCZD17


Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 31-310 of human NRXN1 (NP_001129131.1).

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images



Western blot analysis of extracts of various cell lines, using NRXN1 antibody (CAB10066) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.