
## **CD53 Rabbit Polyclonal Antibody**

## CAB10079



Product Information Size: 20uL, 50uL, 100uL, 200uL	<b>Protein Background</b> The protein encoded by this gene is a member of the transmembrane 4 superfamily, also know as the tetraspanin family. Most of these members are cell-surface proteins that ar characterized by the presence of four hydrophobic domains. The proteins mediate signa		
		Observed MW:	transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with
		35kDa Calculated MW: 24kDa Applications:	integrins. It contributes to the transduction of CD2-generated signals in T cells and natural kille cells and has been suggested to play a role in growth regulation. Familial deficiency of thi gene has been linked to an immunodeficiency associated with recurrent infectious disease caused by bacteria, fungi and viruses. Alternative splicing results in multiple transcript variant <b>Immunogen information</b>
WB	Gene ID: 963		
Reactivity:			
Human, Mouse	Uniprot P19397		
Antibody Information	<b>Synonyms:</b> CD53; MOX44; TSPAN25		
Recommended dilutions: WB 1:1000 - 1:4000			
Source:	Immunogen:		
Rabbit	Recombinant fusion protein containing a sequence corresponding to amino acids 100-180 of human CD53 (NP_000551.1).		
lsotype:			
IgG	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.		

Affinity purification



Western blot analysis of extracts of various cell lines, using CD53 antibody (CAB10079) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.