## **CACNA2D3 Rabbit Polyclonal Antibody**



## CAB10315

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

123kDa

Calculated MW:

59kDa/112kDa/123kDa

**Applications:** 

Reactivity:

WB

Human, Mouse, Rat

**Antibody Information** 

**Recommended dilutions:** 

WB 1:1000 - 1:2000

Source:

Rabbit

Isotype:

IgG

**Purification:** Affinity purification **Protein Background** 

This gene encodes a member of the alpha-2/delta subunit family, a protein in the voltagedependent calcium channel complex. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization and consist of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. Various versions of each of these subunits exist, either expressed from similar genes or the result of alternative splicing. Research on a highly similar protein in rabbit suggests the protein described in this record is cleaved into alpha-2 and delta subunits. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized.

Immunogen information

Gene ID:

55799

Uniprot Q8IZS8

Synonyms:

CACNA2D3; HSA272268

Immunogen:


Recombinant fusion protein containing a sequence corresponding to amino acids 900-1040 of human CACNA2D3 (NP\_060868.2).

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extracts of various cell lines, using CACNA2D3 antibody (CAB10315) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.