

NFKB1 Rabbit Polyclonal Antibody



CAB11160

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

120kDa

Calculated MW:

85kDa/105kDa

Applications:

WB IHC IF IP

Reactivity:

Human

Antibody Information

Recommended dilutions:

WB 1:500 - 1:1000 IHC 1:50
- 1:200 IF 1:20 - 1:100 IP
1:20 - 1:50

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Protein Background

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed.

Immunogen information

Gene ID:

4790

Uniprot

P19838

Synonyms:

NFKB1; CVID12; EBP-1; KBF1; NF-kB1; NF-kappa-B; NF-kappaB;
NFKB-p105; NFKB-p50; NFkappaB; p105; p50


Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 740-964 of human NFKB1 (NP_001158884.1).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images




Western blot analysis of extracts of various cell lines, using NFKB1 antibody (CAB11160) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.



Immunohistochemistry of paraffin-embedded human tonsil using NFKB1 antibody (CAB11160) at dilution of 1:200 (40x lens).



Immunofluorescence analysis of MCF7 cells using NFKB1 antibody (CAB11160).



Immunoprecipitation analysis of 150ug extracts of MCF-7 cells using 3ug NFKB1 antibody (CAB11160). Western blot was performed from the immunoprecipitate using NFKB1 antibody (CAB11160) at a dilution of 1:500.