

# APEX1 Rabbit Polyclonal Antibody



CAB1117

---

## Product Information

### Size:

20uL, 50uL, 100uL, 200uL

### Observed MW:

39kDa

### Calculated MW:

35kDa

### Applications:

WB IHC IF

### Reactivity:

Human

## Protein Background

Apurinic/aprimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that remove specific abnormal bases. AP sites are pre-mutagenic lesions that can prevent normal DNA replication so the cell contains systems to identify and repair such sites. Class II AP endonucleases cleave the phosphodiester backbone 5' to the AP site. This gene encodes the major AP endonuclease in human cells. Splice variants have been found for this gene; all encode the same protein.

## Immunogen information

### Gene ID:

328

### Uniprot

P27695

### Synonyms:

APEX1; APE; APE1; APEN; APEX; APX; HAP1; REF1

## Antibody Information

### Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50  
- 1:100 IF 1:50 - 1:100

### Source:

Rabbit

### Isotype:

IgG

### Purification:

Affinity purification

### Immunogen:


Recombinant fusion protein containing a sequence corresponding to amino acids 1-318 of human APEX1 (NP\_542380.1).

### Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images


---




Western blot analysis of extracts of various cell lines, using APEX1 antibody (CAB1117). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunohistochemistry of paraffin-embedded human liver damage using APEX1 antibody (CAB1117) at dilution of 1:100 (40x lens).



Immunofluorescence analysis of HeLa cells using APEX1 antibody (CAB1117). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of A549 cells using APEX1 antibody (CAB1117).