
Parkin Rabbit Polyclonal Antibody

CAB11172



roduct Information	Protein Background
Size:	The precise function of this gene is unknown; however, the encoded protein is a component of
20uL, 50uL, 100uL, 200uL	a multiprotein E3 ubiquitin ligase complex that mediates the targeting of substrate proteins for proteasomal degradation. Mutations in this gene are known to cause Parkinson disease and
Observed MW:	autosomal recessive juvenile Parkinson disease. Alternative splicing of this gene produces multiple transcript variants encoding distinct isoforms. Additional splice variants of this gene
52kDa	have been described but currently lack transcript support.
Calculated MW:	Immunogen information
23kDa/30-51kDa	Gene ID:
Applications:	5071
WB IF	Uniprot
Reactivity:	O60260
Human, Mouse, Rat	Synonyms:
	PRKN; AR-JP; LPRS2; PARK2; PDJ; Parkin
Antibody Information	
Recommended dilutions:	Immunogen:
WB 1:500 - 1:2000 IF 1:50 -	Recombinant fusion protein containing a sequence corresponding
1:200	to amino acids 1-300 of human Parkin
Source:	
Rabbit	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
lsotype:	sodium azide, 50% glycerol, pH7.3.
lgG	

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using Parkin antibody (CAB11172) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 5s.

Immunofluorescence analysis of A549 cells using Parkin antibody (CAB11172).