
Product Information

Product SKU:	CAB11337	Gene ID:	1029	Size:	20uL, 100uL
Clone No:	-	Host Species:	Rabbit	Reactivity:	Human

Additional Information

Observed MW:	16kDa	Conjugate:	Unconjugated
Calculated MW:	8kDa/11kDa/12kDa/13kDa/16kDa/17kDa	Isotype:	IgG

Immunogen Information

Background:	This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.
Recommended Dilution:	WB, 1:500 - 1:1000
Synonyms:	ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A; CDKN2A/p16INK4a
Purification Method:	Affinity purification
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-156 of human CDKN2A/p16INK4a (NP_000068.1).
Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.