## **IL1 beta Rabbit Polyclonal Antibody**



## **CAB11369**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

17kDa/30kDa

Calculated MW:

30kDa

**Applications:** 

WB

Reactivity:

Human, Mouse, Rat

**Protein Background** 

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

Immunogen information

Gene ID: 3553

Uniprot

P01584

Synonyms: IL-1; IL1-BETA; IL1F2; IL1 beta; IL1B

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Immunogen:

Recombinant fusion protein containing a sequence corresponding Source:

to amino acids 124-223 of human IL1 beta (NP\_000567.1). Rabbit

Storage: Isotype:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% IgG

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using IL1 beta antibody (CAB11369) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.