## **LDHA Rabbit Polyclonal Antibody**



## **CAB1146**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

37kDa

**Calculated MW:** 

26kDa/30kDa/36kDa/39kDa

Applications:

WB IF IP

Reactivity:

Human, Mouse, Rat

**Protein Background** 

The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. The protein is found predominantly in muscle tissue and belongs to the lactate dehydrogenase family. Mutations in this gene have been linked to exertional myoglobinuria. Multiple transcript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene.

Immunogen information

Gene ID:

3939

Uniprot P00338

Synonyms:

GSD11; HEL-S-133P; LDHM; PIG19; LDHA

**Antibody Information** 

Recommended dilutions:

WB 1:500 - 1:2000 IF 1:50 - 1:200 IP 1:50 - 1:100

Source:

Rabbit

Isotype:

lgG

Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 1-332 of human LDHA (NP\_005557.1).

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.


**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using LDHA antibody (CAB1146) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of A549 cells using LDHA antibody (CAB1146). Blue: DAPI for nuclear staining.