
MC3R Rabbit Polyclonal Antibody

CAB11609



| roduct Information | Protein Background |
|--|--|
| Size: | This gene encodes a G-protein-coupled receptor for melanocyte-stimulating hormone and |
| 20uL, 50uL, 100uL, 200uL | adrenocorticotropic hormone that is expressed in tissues other than the adrenal cortex and melanocytes. This gene maps to the same region as the locus for benign neonatal epilepsy |
| Observed MW: | Mice deficient for this gene have increased fat mass despite decreased food intake, suggesting a role for this gene product in the regulation of energy homeostasis. Mutations in this gene are |
| 36kDa | associated with a susceptibility to obesity in humans. |
| Calculated MW: | Immunogen information |
| 36kDa | Gene ID: |
| Applications: | 4159 |
| WB | Uniprot |
| Reactivity: | P41968 |
| Mouse, Rat | Synonyms: MC3R; BMIQ9; MC3; MC3-R; OB20; OQTL |
| Antibody Information | |
| Recommended dilutions: WB 1:500 - 1:2000 | Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human MC3R (NP_063941.3). |
| Source: Rabbit | |
| | Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% |
| lsotype: lgG | sodium azide, 50% glycerol, pH7.3. |

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using MC3R antibody (CAB11609) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.