## **MMP13 Rabbit Polyclonal Antibody**



## **CAB11755**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

43KDa/54KDa/68KDa

**Calculated MW:** 

53kDa

**Applications:** 

WB, IF

Reactivity:

Human, Mouse, Rat

Category:

**Polyclonal Antibodies** 

**Protein Background** 

This gene encodes a member of the peptidase M10 family of matrix metalloproteinases (MMPs). Proteins in this family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. The encoded preproprotein is proteolytically processed to generate the mature protease. This protease cleaves type II collagen more efficiently than types I and III. It may be involved in articular cartilage turnover and cartilage pathophysiology associated with osteoarthritis. Mutations in this gene are associated with metaphyseal anadysplasia. This gene is part of a cluster of MMP genes on chromosome 11.

Immunogen information

**Gene ID:** 4322

**Uniprot** P45452

Synonyms:

MMP13; CLG3; MANDP1; MDST; MMP-13

**Antibody Information** 

Recommended dilutions:

WB 1:500 - 1:2000 IF 1:50 - 1:200

Source:

Isotype:

Rabbit

DDIC

lgG

Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 262-471 of human MMP13 (NP\_002418.1).

Storage:


Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**





Western blot analysis of extracts of various cell lines, using MMP13 Rabbit pAb (CAB11755) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% non-fat dry milk in TBST.

Detection: ECL Basic Kit (AbGn00020).

Exposure time: 3s.

Immunofluorescence analysis of L929 cells using MMP13 Rabbit pAb (CAB11755) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.