
KRT7 Rabbit Polyclonal Antibody

CAB12004



roduct Information	Protein Background
Size:	The protein encoded by this gene is a member of the keratin gene family. The type I
20uL, 50uL, 100uL, 200uL	cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This
Observed MW:	type II cytokeratin is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels. The genes encoding the type I
Refer to figures	cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing ma result in several transcript variants; however, not all variants have been fully described.
Calculated MW:	result in several transcript variants, new ever, not an variants have been rany described.
51kDa	Immunogen information
Applications:	Gene ID: 3855
WB IHC IF	
Reactivity:	Uniprot P08729
Human, Mouse, Rat	
	Synonyms: KRT7; CK7; K2C7; K7; SCL; keratin 7; Keratin
Antibody Information	
Recommended dilutions: WB 1:500 - 1:2000 IHC 1:50	Immunogen:
- 1:200 IF 1:50 - 1:200	Recombinant fusion protein containing a sequence corresponding
Source: Rabbit	to amino acids 1-94 of human KRT7 (NP_005547.3).
	Storage:
lsotype: lgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Immunohistochemistry of paraffin-embedded human lung cancer using KRT7 antibody (CAB12004) at dilution of 1:200 (40x lens).

Immunofluorescence analysis of human lung cancer using KRT7 antibody (CAB12004) at dilution of 1:100. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of mouse bronchus using KRT7 antibody (CAB12004) at dilution of 1:100. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of rat bronchus using KRT7 antibody (CAB12004) at dilution of 1:100. Blue: DAPI for nuclear staining.