
## **LTBR Rabbit Polyclonal Antibody**

## CAB12006



roduct Information	Protein Background
Size:	This gene encodes a member of the tumor necrosis factor receptor superfamily. The majo
20uL, 50uL, 100uL, 200uL	ligands of this receptor include lymphotoxin alpha/beta and tumor necrosis factor ligand superfamily member 14. The encoded protein plays a role in signalling during the developmen
Observed MW:	of lymphoid and other organs, lipid metabolism, immune response, and programmed ce death. Activity of this receptor has also been linked to carcinogenesis. Alternatively splice
47kDa	transcript variants encoding multiple isoforms have been observed.
Calculated MW:	Immunogen information
44kDa/46kDa	Gene ID:
Applications:	4055
WB	Uniprot
Reactivity:	P36941
Mouse	Synonyms:
	LTBR; CD18; D12S370; LT-BETA-R; TNF-R-III; TNFCR; TNFR-RP; TNFR2-RP; TNFR3; TNFRSF3
Antibody Information	
Recommended dilutions:	Immunogen:
WB 1:500 - 1:2000	Recombinant fusion protein containing a sequence corresponding to amino acids 249-435 of human LTBR (NP_002333.1).
<b>Source:</b> Rabbit	
Ναυσιτ	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
<b>lsotype:</b> lgG	sodium azide, 50% glycerol, pH7.3.

**Purification:** Affinity purification



Western blot analysis of extracts of mouse liver, using LTBR Antibody (CAB12006) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.