## **PORCN Rabbit Polyclonal Antibody**



## **CAB12120**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

60kDa

Calculated MW:

42kDa/51kDa/52kDa

**Applications:** 

WB

Reactivity:

Human

**Protein Background** 

This gene belongs to the evolutionarily conserved porcupine (Porc) gene family. Genes of the porcupine family encode endoplasmic reticulum proteins with multiple transmembrane domains. Porcupine proteins are involved in the processing of Wnt (wingless and int homologue) proteins. Disruption of this gene is associated with focal dermal hypoplasia, and the encoded protein has been implicated in cancer. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed.

Immunogen information

Gene ID:

64840

Uniprot Q9H237

**Synonyms:** 

PORCN; DHOF; FODH; MG61; PORC; PPN

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

Immunogen:

Recombinant fusion protein containing a sequence corresponding

to amino acids 220-320 of human PORCN (NP\_982299.1).

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using PORCN antibody (CAB12120) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 60s.