COASY Rabbit Polyclonal Antibody



CAB12179

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

62kDa

Calculated MW:

62kDa/65kDa

Applications:

WB IHC

Reactivity:

Human, Mouse, Rat

Protein Background

Coenzyme A (CoA) functions as a carrier of acetyl and acyl groups in cells and thus plays an important role in numerous synthetic and degradative metabolic pathways in all organisms. In eukaryotes, CoA and its derivatives are also involved in membrane trafficking and signal transduction. This gene encodes the bifunctional protein coenzyme A synthase (CoAsy) which carries out the last two steps in the biosynthesis of CoA from pantothenic acid (vitamin B5). The phosphopantetheine adenylyltransferase domain of this bifunctional protein catalyzes the conversion of 4'-phosphopantetheine into dephospho-coenzyme A (dpCoA) while its dephospho-CoA kinase domain completes the final step by phosphorylating dpCoA to form CoA. Mutations in this gene are associated with neurodegeneration with brain iron accumulation (NBIA). Alternative splicing results in multiple isoforms.

Immunogen information

Gene ID: 80347

Uniprot Q13057

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:100 - 1:200

Source:

Rabbit

Isotype: IgG

Synonyms:

COASY; DPCK; NBIA6; NBP; PPAT; UKR1; pOV-2

Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 345-564 of human COASY (NP_079509.5).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification

Product Images



Western blot analysis of extracts of various cell lines, using COASY antibody (CAB12179) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.