## **PPIH Rabbit Polyclonal Antibody**



## **CAB12274**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

19kDa

**Calculated MW:** 

14kDa/19kDa

**Applications:** 

WB

Reactivity:

Human, Rat

?:\_\_.

**Protein Background** 

The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPlase) family. PPlases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is a specific component of the complex that includes pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. This protein has been shown to possess PPlase activity and may act as a protein chaperone that mediates the interactions between different proteins inside the spliceosome.

Immunogen information

**Gene ID:** 10465

**Uniprot** O43447

Synonyms:

PPIH; CYP-20; CYPH; SnuCyp-20; USA-CYP

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Immunogen:

Recombinant fusion protein containing a sequence corresponding

**Source:** to amino acids 1-177 of human PPIH (NP\_006338.1).

Rabbit

Storage


**Isotype:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

lgG sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using PPIH antibody (CAB12274) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 60s.