SURF1 Rabbit Polyclonal Antibody


CAB12521



Product Information	Protein Background
Size:	This gene encodes a protein localized to the inner mitochondrial membrane and thought to be
20uL, 50uL, 100uL, 200uL	involved in the biogenesis of the cytochrome c oxidase complex. The protein is a member of the SURF1 family, which includes the related yeast protein SHY1 and rickettsial protein RP733.
Observed MW:	The gene is located in the surfeit gene cluster, a group of very tightly linked genes that do not share sequence similarity, where it shares a bidirectional promoter with SURF2 on the opposite
33kDa	strand. Defects in this gene are a cause of Leigh syndrome, a severe neurological disorder that is commonly associated with systemic cytochrome c oxidase deficiency.
Calculated MW:	is commonly associated with systemic cytochrome c oxidase deficiency.
31kDa/33kDa	Immunogen information
Applications:	Gene ID: 6834
WB IHC	
Reactivity:	Uniprot Q15526
Human, Mouse, Rat	
	Synonyms: SURF1; CMT4K
Antibody Information	
Recommended dilutions: WB 1:500 - 1:2000 IHC 1:50 - 1:200	Immunogen: Recombinant fusion protein containing a sequence corresponding
Source: Rabbit	to amino acids 80-275 of human SURF1 (NP_003163.1).
	Storage:
lsotype: IgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification

www.assaygenie.com



Western blot analysis of extracts of various cell lines, using SURF1 antibody (CAB12521) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 3s.

Immunohistochemistry of paraffin-embedded rat kidney using SURF1 antibody (CAB12521) at dilution of 1:100 (40x lens).