HIRA Rabbit Polyclonal Antibody



CAB12527

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

112kDa

Calculated MW:

89kDa/111kDa

Applications:

WB

Reactivity:

Human, Mouse

Uniprot P54198

Gene ID:

7290

Synonyms:

HIRA; DGCR1; TUP1; TUPLE1

Immunogen information

Protein Background

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source: Rabbit

Isotype:

IgG

Immunogen:

Recombinant fusion protein containing a sequence corresponding

This gene encodes a histone chaperone that preferentially places the variant histone H3.3 in nucleosomes. Orthologs of this gene in yeast, flies, and plants are necessary for the formation

of transcriptionally silent heterochomatin. This gene plays an important role in the formation of the senescence-associated heterochromatin foci. These foci likely mediate the irreversible

cell cycle changes that occur in senescent cells. It is considered the primary candidate gene in some haploinsufficiency syndromes such as DiGeorge syndrome, and insufficient production

to amino acids 250-550 of human HIRA (NP_003316.3).

of the gene may disrupt normal embryonic development.


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification

Product Images



Western blot analysis of extracts of various cell lines, using HIRA Antibody (CAB12527) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.