
CHRFAM7A Rabbit Polyclonal Antibody

CAB12603



Product Information Size: 20uL, 50uL, 100uL, 200uL Observed MW: 50kDa Calculated MW: 46kDa Applications:	Protein Background The nicotinic acetylcholine receptors (nAChRs) are members of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. The family member CHRNA7, which is located on chromosome 15 in a region associated with several neuropsychiatric disorders, is partially duplicated and forms a hybrid with a novel gene from the family with sequence similarity 7 (FAM7A). Alternative splicing has been observed, and two variants exist, for this hybrid gene. The N-terminally truncated products predicted by the largest open reading frames for each variant would lack the majority of the neurotransmitter-gated ion-channel ligand binding domain but retain the transmembrane region that forms the ion channel. Although current evidence supports transcription of this hybrid gene, translation of the nicotinic acetylcholine receptor-like protein-encoding open reading frames has not been confirmed.
WB	Immunogen information
Reactivity: Human, Mouse, Rat	Gene ID: 89832 Uniprot
	Q494W8
Antibody Information	
Recommended dilutions: WB 1:500 - 1:2000	Synonyms: CHRFAM7A; CHRNA7; CHRNA7-DR1; D-10
Source: Rabbit Isotype: IgG	Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human CHRFAM7A (NP_647536.1).
Purification: Affinity purification	Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



Western blot analysis of extracts of various cell lines, using CHRFAM7A antibody (CAB12603) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.