NDUFS8 Rabbit Polyclonal Antibody

CAB13034




Product Information	Protein Background
Size:	This gene encodes a subunit of mitochondrial NADH:ubiquinone oxidoreductase, or Complex
20uL, 50uL, 100uL, 200uL	I, a multimeric enzyme of the respiratory chain responsible for NADH oxidation, ubiquinone reduction, and the ejection of protons from mitochondria. The encoded protein is involved in
Observed MW:	the binding of two of the six to eight iron-sulfur clusters of Complex I and, as such, is required in the electron transfer process. Mutations in this gene have been associated with Leigh
23kDa	syndrome.
Calculated MW:	Immunogen information
23kDa	Gene ID:
Applications:	4728
WB IHC IF	Uniprot
Reactivity:	O00217
Human, Mouse, Rat	Synonyms: NDUFS8; CI-23k; CI23KD; TYKY
Antibody Information	
Recommended dilutions:	Immunogen:
WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:50 - 1:100	Recombinant fusion protein containing a sequence corresponding to amino acids 35-210 of human NDUFS8 (NP_002487.1).
Source: Rabbit	
	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%


sodium azide, 50% glycerol, pH7.3.

lsotype: lgG

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using NDUFS8 antibody (CAB13034) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.



Immunofluorescence analysis of L929 cells using NDUFS8 Polyclonal Antibody (CAB13034) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.