

Slug Rabbit Polyclonal Antibody



CAB13352

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

30KDa

Calculated MW:

29kDa

Applications:

WB IHC

Reactivity:

Human, Mouse, Rat

Protein Background

This gene encodes a member of the Snail family of C2H2-type zinc finger transcription factors. The encoded protein acts as a transcriptional repressor that binds to E-box motifs and is also likely to repress E-cadherin transcription in breast carcinoma. This protein is involved in epithelial-mesenchymal transitions and has antiapoptotic activity. Mutations in this gene may be associated with sporadic cases of neural tube defects.

Immunogen information

Gene ID:

6591

Uniprot

O43623

Synonyms:

SNAI2; SLUG; SLUGH; SLUGH1; SNAIL2; WS2D

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50
- 1:200

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification


Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 126-268 of human Slug (NP_003059.1).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images




Western blot analysis of extracts of various cell lines, using Slug antibody (CAB13352) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 60s.



Immunohistochemistry of paraffin-embedded human liver damage using Slug Antibody (CAB13352) at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human breast cancer using Slug Antibody (CAB13352) at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human stomach using Slug Antibody (CAB13352) at dilution of 1:100 (40x lens).