

# HDAC4 Rabbit Monoclonal Antibody



CAB13510

## Product Information

### Size:

20uL, 50uL, 100uL, 200uL

### Observed MW:

140KDa

### Calculated MW:

140kDa

### Applications:

WB

### Reactivity:

Mouse, Rat

## Antibody Information

### Recommended dilutions:

WB 1:500 - 1:2000

### Source:

Rabbit

### Isotype:

IgG

### Purification:

Affinity purification

## Protein Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3. [provided by RefSeq, Jul 2008]

## Immunogen information

### Gene ID:

9759

### Uniprot

P56524

### Synonyms:

AHO3; BDMR; HA6116; HD4; HDAC-4; HDAC-A; HDACA

### Immunogen:


A synthesized peptide derived from human HDAC4

### Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Product Images

---



Western blot - HDAC4 Rabbit mAb (CAB13510)