## **TSPAN7 Rabbit Polyclonal Antibody**



## **CAB13555**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

35KDa

**Calculated MW:** 

27kDa

Applications:

WB

Reactivity:

Human

**Protein Background** 

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein and may have a role in the control of neurite outgrowth. It is known to complex with integrins. This gene is associated with X-linked mental retardation and neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome and myotonic dystrophy.

Immunogen information

**Gene ID:** 7102

**Uniprot** P41732

Synonyms:

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

VVD 1.300 - 1.2000

Source:

Rabbit

Immunogen:

TM4SF2; TM4SF2b

Recombinant fusion protein containing a sequence corresponding to amino acids 113-213 of human TSPAN7 (NP\_004606.2).

TSPAN7; A15; CCG-B7; CD231; DXS1692E; MRX58; MXS1; TALLA-1;

Isotype: Storage:


IgG Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of Jurkat cells, using TSPAN7 antibody (CAB13555) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 15s.