## **RPLP0 Rabbit Polyclonal Antibody**





**Product Information** 

**Product SKU**: CAB13633 **Gene ID**: 6175 **Size**: 20uL, 100uL

Clone No: - Host Species: Rabbit Reactivity: Human, Mouse

**Additional Information** 

**Observed MW**: 37kDa **Conjugate:** Unconjugated

Calculated MW: 34kDa Isotype: IgG

## Immunogen Information

**Background**: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S

subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed

pseudogenes of this gene dispersed through the genome.

**Recommended Dilution**: WB,1:500 - 1:2000 IF/ICC,1:50 - 1:200

Synonyms: P0; LP0; L10E; RPP0; uL10; PRLP0; RPLP0

**Purifcation Method**: Affinity purification

**Immunogen**: Recombinant fusion protein containing a sequence corresponding to amino acids 1-317 of human

RPLP0 (NP\_000993.1).

**Storage**: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.