## **FN3K Rabbit Polyclonal Antibody**



## **CAB13727**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

35kDa

Calculated MW:

35kDa

Applications:

WB

Reactivity:

Human, Mouse

**Protein Background** 

A high concentration of glucose can result in non-enzymatic oxidation of proteins by reaction of glucose and lysine residues (glycation). Proteins modified in this way, fructosamines, are less active or functional. This gene encodes an enzyme which catalyzes the phosphorylation of fructosamines which may result in deglycation.

Immunogen information

**Gene ID:** 64122

**Uniprot** Q9H479

**Synonyms:** 

Immunogen:

FN3K

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source:

Rabbit

**Isotype:** IgG Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

Recombinant fusion protein containing a sequence corresponding


to amino acids 1-200 of human FN3K (NP\_071441.1).

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using FN3K antibody (CAB13727) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.