
CD74 Rabbit Polyclonal Antibody

CAB13958



roduct Information	Protein Background
Size:	The protein encoded by this gene associates with class II major histocompatibility complex
20uL, 50uL, 100uL, 200uL	(MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration
Observed MW:	inhibitory factor (MIF) which, when bound to the encoded protein, initiates survival pathway and cell proliferation. This protein also interacts with amyloid precursor protein (APP) and
30-40kDa	suppresses the production of amyloid beta (Abeta). Multiple alternatively spliced transcrip variants encoding different isoforms have been identified.
Calculated MW:	
18kDa/26kDa/33kDa	Immunogen information
Applications:	Gene ID: 972
WB	
Reactivity:	Uniprot P04233
Human, Mouse	
	Synonyms: CD74; DHLAG; HLADG; II; Ia-GAMMA
Antibody Information	
Recommended dilutions:	
WB 1:500 - 1:2000	Immunogen:
_	Recombinant fusion protein containing a sequence corresponding to amino acids 73-232 of human CD74 (NP_004346.1).
Source: Rabbit	
	Storage:
lsotype: IgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using CD74 antibody (CAB13958) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 60s.