KCNN3 Rabbit Polyclonal Antibody

CAB14012



Product Information Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

50-85kDa

Calculated MW:

47kDa/48kDa/82kDa

Applications:

WB IF

Reactivity:

Human, Mouse, Rat

Antibody Information

–

Recommended dilutions: WB 1:500 - 1:1000 IF 1:50 -

WB 1:500 - 1:1000 IF 1:50 1:100

Source: Rabbit

Immunogen:

Recombinant fusion protein containing a sequence corresponding to amino acids 237-426 of human KCNN3 (NP_740752.1).

Isotype: IgG

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification

Protein Background


Action potentials in vertebrate neurons are followed by an afterhyperpolarization (AHP) that may persist for several seconds and may have profound consequences for the firing pattern of the neuron. Each component of the AHP is kinetically distinct and is mediated by different calcium-activated potassium channels. This gene belongs to the KCNN family of potassium channels. It encodes an integral membrane protein that forms a voltage-independent calciumactivated channel, which is thought to regulate neuronal excitability by contributing to the slow component of synaptic AHP. This gene contains two CAG repeat regions in the coding sequence. It was thought that expansion of one or both of these repeats could lead to an increased susceptibility to schizophrenia or bipolar disorder, but studies indicate that this is probably not the case. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Immunogen information

Gene ID: 3782

Uniprot Q9UGI6

Synonyms: KCNN3; KCa2.3; SK3; SKCA3; hSK3



Western blot analysis of extracts of various cell lines, using KCNN3 antibody (CAB14012) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (CABM00021). Exposure time: 60s.

Immunofluorescence analysis of U2OS cells using KCNN3 antibody (CAB14012). Blue: DAPI for nuclear staining.