
SIRT2 Rabbit Polyclonal Antibody

CAB14109



roduct Information	Protein Background
Size:	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2
20uL, 50uL, 100uL, 200uL	protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeas
Observed MW:	sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory
39kDa	proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative
Calculated MW:	splicing of this gene.
30kDa/35kDa/39kDa/41kDa/ 43kDa	Immunogen information
Applications:	Gene ID: 22933
WB	
Reactivity:	Uniprot Q8IXJ6
Mouse, Rat	
	Synonyms: SIRT2; SIR2; SIR2L; SIR2L2; sirtuin 2
Antibody Information	
Recommended dilutions:	
WB 1:500 - 1:2000	Immunogen:
	A synthetic peptide of human SIRT2
Source: Rabbit	
	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
lsotype: lgG	sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using SIRT2 antibody (CAB14109) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.