
Cyclin A1 Rabbit Polyclonal Antibody

CAB14529



roduct Information	Protein Background
Size:	The protein encoded by this gene belongs to the highly conserved cyclin family, whose
20uL, 50uL, 100uL, 200uL	members are characterized by a dramatic periodicity in protein abundance through the cel cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression
Observed MW:	and degradation patterns which contribute to the temporal coordination of each mitotic event The cyclin encoded by this gene was shown to be expressed in testis and brain, as well as ir
52kDa	several leukemic cell lines, and is thought to primarily function in the control of the germline meiotic cell cycle. This cyclin binds both CDK2 and CDC2 kinases, which give two distinct kinase
Calculated MW:	activities, one appearing in S phase, the other in G2, and thus regulate separate functions ir
47kDa/52kDa	cell cycle. This cyclin was found to bind to important cell cycle regulators, such as Rb family proteins, transcription factor E2F-1, and the p21 family proteins. Multiple transcript variants
Applications:	encoding different isoforms have been found for this gene.
WB	Immunogen information
Reactivity:	Gene ID:
Mouse, Rat	8900
	Uniprot P78396
Antibody Information	
Recommended dilutions: WB 1:500 - 1:2000	Synonyms: CCNA1; CT146; cyclin-A1
Source: Rabbit	
Ναυσιι	Immunogen:
	A synthetic peptide corresponding to a sequence within amino
lsotype: IgG	acids 350-450 of human CCNA1 (NP_001104515.1).
	Storage:
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%
Purification:	sodium azide, 50% glycerol, pH7.3.
Purification: Affinity purification	



Western blot analysis of extracts of various cell lines, using Cyclin A1 antibody (CAB14529) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (CABS003) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 10s.