PDP1 Rabbit Polyclonal Antibody





Product Information

Product SKU: CAB14545 **Gene ID**: 54704 **Size**: 20uL, 100uL

Clone No: - Host Species: Rabbit Reactivity: Human, Mouse, Rat

Additional Information

Observed MW: 61kDa **Conjugate:** Unconjugated

Calculated MW: 61kDa Isotype: IgG

Immunogen Information

Background: Pyruvate dehydrogenase (E1) is one of the three components (E1, E2, and E3) of the large pyruvate

dehydrogenase complex. Pyruvate dehydrogenase kinases catalyze phosphorylation of serine residues of E1 to inactivate the E1 component and inhibit the complex. Pyruvate dehydrogenase phosphatases catalyze the dephosphorylation and activation of the E1 component to reverse the effects of pyruvate dehydrogenase kinases. Pyruvate dehydrogenase phosphatase is a heterodimer consisting of catalytic and regulatory subunits. Two catalytic subunits have been reported; one is predominantly expressed in skeletal muscle and another one is is much more abundant in the liver. The catalytic subunit, encoded by this gene, is the former, and belongs to the protein phosphatase 2C (PP2C) superfamily. Along with the pyruvate dehydrogenase complex and pyruvate dehydrogenase kinases, this enzyme is located in the mitochondrial matrix. Mutation in this gene causes pyruvate dehydrogenase phosphatase deficiency.

Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Recommended Dilution: WB,1:500 - 1:2000 IF/ICC,1:50 - 1:200

Synonyms: PDH; PDP; PDPC; PPM2A; PPM2C; PDP1

Purifcation Method: Affinity purification

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 72-260 of human

PDP1 (NP_060914.2).

Storage: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.