
## HABP2 Rabbit Polyclonal Antibody

## CAB14565



roduct Information Size:	<b>Protein Background</b> This gene encodes a member of the peptidase S1 family of serine proteases. The encode
	preproprotein is secreted by hepatocytes and proteolytically processed to generate heavy and
20uL, 50uL, 100uL, 200uL	light chains that form the mature heterodimer. Further autoproteolysis leads to smaller, inactive peptides. This extracellular protease binds hyaluronic acid and may play a role in the
Observed MW:	coagulation and fibrinolysis systems. Mutations in this gene are associated with nonmedullar thyroid cancer and susceptibility to venous thromboembolism. Alternative splicing results in
63kDa	multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically
Calculated MW:	processed.
59kDa/62kDa	Immunogen information
Applications:	Gene ID:
WB	3026
Reactivity:	Uniprot
Human, Mouse, Rat	Q14520
Antibody Information	<b>Synonyms:</b> HABP2; FSAP; HABP; HGFAL; NMTC5; PHBP
-	
Recommended dilutions: WB 1:500 - 1:2000	
	Immunogen:
<b>Source:</b> Rabbit	Recombinant fusion protein containing a sequence corresponding to amino acids 350-560 of human HABP2 (NP_004123.1).
lsotype:	Storage:
lgG	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purification:** Affinity purification



Western blot analysis of extracts of various cell lines, using HABP2 antibody (CAB14565) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.