
Product Information

Product SKU:	CAB1483	Gene ID:	1026	Size:	20uL, 100uL
Clone No:	-	Host Species:	Rabbit	Reactivity:	Human,Mouse,Rat

Additional Information

Observed MW:	18kDa(Mouse)/21kDa(Human/Rat)	Conjugate:	Unconjugated
Calculated MW:	18kDa	Isotype:	IgG

Immunogen Information

Background:	This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.
Recommended Dilution:	WB,1:500 - 1:1000 IF/ICC,1:50 - 1:200 IHC-P,1:50 - 1:200
Synonyms:	P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1; CDKN1A/p21CIP1
Purification Method:	Affinity purification
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 1-164 of human CDKN1A/p21CIP1 (NP_000380.1).
Storage:	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.